

A remark on a question of Beauville about lagrangian fibrations

Ekaterina Amerik*

Abstract

This note is a proof of the fact that a lagrangian torus on an irreducible hyperkähler fourfold is always a fiber of an almost holomorphic lagrangian fibration.

AMS 2010 Subject Classification: 14J99, 32J27

Keywords: holomorphic symplectic manifolds, lagrangian tori, fibrations.

Let X be an irreducible holomorphic symplectic variety of dimension $2n$, with a lagrangian torus $A \subset X$ (such a torus is always an abelian variety, even if X is not projective: this follows from the fact that the space $H^{2,0}(X)$ restricts to zero on A , see [C]). Beauville [B] asked whether there always exists a lagrangian fibration f (holomorphic or almost holomorphic), such that A is a fiber of f . Recently, D. Greb, C. Lehn and S. Rollenske [GLR] proved that it is always the case when X is non-projective, or even admits a non-projective deformation preserving the lagrangian subtorus. It is not clear a priori whether all irreducible holomorphic symplectic varieties admit such deformations. The authors also have announced another paper in preparation, where Beauville's question is answered in the affirmative in dimension four¹. The purpose of this short note is to provide an answer to the "almost holomorphic version" of Beauville's question in dimension four in a very elementary way.

Since the torus $A \subset X$ is lagrangian, its normal bundle $N_{A,X}$, being isomorphic to its cotangent bundle, is trivial. The deformation theory of lagrangian tori in holomorphic symplectic manifolds is well understood: as it is explained in [GLR],

*Université Paris-Sud, Campus d'Orsay, Bât. 425, 91405 Orsay, France; and National Research University Higher School of Economics, Department of Mathematics, Vavilova 7, 117332, Moscow, Russia; Ekaterina.Amerik@math.u-psud.fr; the author is partially supported by AG Laboratory, RF government grant ag. 11.G34.31.0023.

¹Their paper appeared on arxiv.org:1110.2680 a few days after a conversation between myself and one of the authors, and contains a simplified (with respect to the original approach of the authors) argument for the existence of a meromorphic fibration, similar to the one given below. My primary reason for making this note public was to provide a source which they could cite.

by the results of Ran and Voisin the space T of deformations of A in X is of dimension n , it is smooth at A , and any deformation A_t of A which is smooth is itself a lagrangian torus. Moreover, if Y denotes the universal family of the A_t 's, the projection $q : Y \rightarrow X$ is unramified along each smooth A_t . In particular, one has a finite number d of A_t 's through a general point of X , and if through a certain point $x \in X$ there is an infinite number of A_t then almost all of them are singular (in fact, if Y denotes the universal family, the projection $q : Y \rightarrow X$ is unramified along each smooth A_t). To show that there is an almost holomorphic fibration with fibers A_t is the same as to show that $d = 1$, or that a general A_t does not intersect any other member A_s .

To say the same things using slightly different words, from the above-mentioned facts on the universal family map q the following is immediate.

Fact: *Locally in a neighbourhood U of A , one has a lagrangian fibration by its small deformations A_t .*

One has to show that this local fibration gives rise to a global, possibly meromorphic, fibration.

Let us list a few immediate consequences of this fact in a lemma. We assume for simplicity that X is projective.

Lemma 1: *a) A lagrangian torus does not intersect its small deformations;
b) Whenever $A_t \cap A_s \neq \emptyset$ and A_s is smooth, all irreducible components of $A_t \cap A_s$ are positive-dimensional;
c) For a general $A_{s'}$ intersecting A_t , the intersection $A_t \cap A_{s'}$ is equidimensional.*

Proof: Whereas a) is immediate from the existence of a local fibration, b) deserves a few words of explanation. Consider the local fibration $f_s : U_s \rightarrow T_s$ in a neighbourhood of A_s (here T_s is a small disc around $s \in T$). The intersection of A_t with U_s may consist of several local components. The statement that $A_t \cap A_s$ cannot have a zero-dimensional component is clearly implied by the claim that no component of $A_t \cap U_s$ dominates the base T_s (indeed, such a component would have zero-dimensional general fiber under the map f_s , whereas for a non-dominating component all fibers are of strictly positive dimension). If this claim does not hold, we can choose a sufficiently general point $s' \in T_s$ outside of the image of all non-dominating components of $A_t \cap U_s$. The resulting lagrangian torus $A_{s'}$ has non-empty zero-dimensional intersection with A_t . This is impossible since the self-intersection number $[A] \cdot [A] = 0$.

The proof of c) is similar to b): it suffices to choose $s' \in T_s$ sufficiently general in a "maximal" image of a component of $A_t \cap U_s$. If this image is of dimension c , the intersection $A_t \cap A_{s'}$ is purely $(n - c)$ -dimensional.

Consider now the case $\dim(X) = 4$, then the intersection $A_t \cap A_s$ is a curve if nonempty. Fix a general lagrangian torus A_t . Since the family of its deformations $A_s, s \in T$ induces a local fibration in the neighbourhood of each smooth member, the intersections $A_t \cap A_s$ induce a fibration or several fibrations on A_t . Indeed,

since most of the A_s intersecting A_t are smooth, this is a family of cycles on A_t not intersecting their neighbours, as are the A_s on X itself; since these cycles are divisors on a surface, those are (possibly reducible) curves whose square is equal to zero, that is, our A_t is fibered in elliptic curves and our intersections are unions of fibers.

Of course, the A_s intersecting A_t do not necessarily form an irreducible family, so apriori there can be several (say, $k > 1$) such fibrations on A_t . Nevertheless an easy linear algebra argument shows that there is in fact only one:

Proposition 2: *One has $k = 1$.*

Proof: Otherwise, we can find two more tori A_u and A_s through a general point p of A_t , in such a way that the pairwise intersections C_{ts} , C_{tu} and C_{us} are curves with distinct tangents at p and the intersection of all three tori has p as an isolated point. One deduces easily from Sard's lemma that for general t, s , the intersection $A_s \cap A_t$ is reduced, so we may assume that the tangent planes to A_u , A_s and A_t are distinct. But since the pairwise intersections of those planes in $T_p X$ are distinct lines, the planes only span a hyperplane $V \subset T_p X$. Now the restriction σ_V of the symplectic form σ_p , that is, the value of σ at p , to V has one-dimensional kernel. Recall that $T_p A_u$, $T_p A_s$ and $T_p A_t$ are σ_p -isotropic. Since $V = T_p A_t + T_p A_s$, one must have $T_p C_{ts} = \text{Ker}(\sigma_V)$. But by the same reason the same holds for $T_p C_{tu}$, a contradiction.

Now we are ready to prove the announced result, which we formulate as a theorem.

Theorem 3: *The answer to the almost holomorphic version of Beauville's question is positive in dimension 4, that is, a general A_t does not intersect any other A_s , so that $d = 1$ and the family A_t induces an almost holomorphic fibration on X .*

Proof: Suppose $d > 1$, then we claim that the irreducible components of the intersections $A_t \cap A_s$ rationally fiber X , that is, there is only one such curve through a general point $p \in X$. Indeed, suppose there are two of them, C_1 and C_2 . Then C_1 is a component of the intersection $A_{1,1} \cap A_{1,2}$ and C_2 of $A_{2,1} \cap A_{2,2}$. Since p is general in X , we may suppose that all four of the above abelian surfaces are smooth. Each pair of those four tori intersect at p , and we know that they should then intersect along a curve through p . Looking at the possible configurations and keeping in mind that the intersection of smooth surfaces is a disjoint union of elliptic curves, one is back to what is just ruled out in Proposition 2.

On the other hand, it is proved in [AC], Proposition 3.4, that any rational fibration $f : X \dashrightarrow B$ of an irreducible holomorphic symplectic variety X has fibers of dimension at least $\dim(X)/2$, provided that the general fiber is not of general type. Since the irreducible components of $A_t \cap A_s$ are elliptic curves, they cannot rationally fiber X , a contradiction.

Remark 4: This argument generalizes to higher-dimensional X in the case when the intersection of two general (intersecting) lagrangian tori $A_t \cap A_s$ is of

codimension one in each; so if $2n = \dim(X)$, one wants $A_t \cap A_s$ to be equal to $n - 1$; when $n = 2$ this is of course the only option unless X is fibered by the A_t 's. More precisely, $A_t \cap A_s$ must be of dimension $n - 1$ for *each* general pair of intersecting lagrangian tori: that is to say, the subset $W = \{(t, s) | A_t \cap A_s \neq \emptyset\} \subset T \times T$ can have several irreducible components W_i , $i = 1, \dots, l$, such that a generic pair of tori in W_i intersects in dimension e_i , and for our argument we need this dimension to be equal to $n - 1$ for any $i = 1, \dots, l$. As soon as this condition is satisfied, the same reasoning as above applies to show the existence of an almost holomorphic lagrangian fibration with fibers A_t .

We hope to return to Beauville's question in a forthcoming joint work with Frédéric Campana.

References

- [AC] E. Amerik, F. Campana: Fibrations méromorphes sur certaines variétés à fibré canonique trivial, Pure Appl. Math. Quarterly 4 no. 2 (special issue in honour of F. Bogomolov), 1-37.
- [B] A. Beauville: Holomorphic symplectic geometry: a problem list, in: Complex and Differential Geometry, Conference held at Leibniz Universitt Hannover (volume 8 of Springer Proceedings in Mathematics), Springer, 2011.
- [C] F. Campana: Isotrivialité de certaines familles kählériennes de variétés non projectives. Math. Z. 252 (2006), 147156.
- [GLR] D. Greb, C. Lehn, S. Rollenske: Lagrangian fibrations on hyperkähler manifolds – on a question of Beauville, arxiv preprint 1105.3410